Conservation Biology 🗞

CONTRIBUTED PAPER

Survival of Eurasian lynx in the human-dominated landscape of Europe

J. Premier ^{1,2,3} M. L. Bastianelli ^{1,2} J. Oeser ⁴ O. Anders ⁵ H. Andren ⁶
M. Aronsson ⁶ 💿 G. Bagrade ⁷ 💿 E. Belotti ^{8,9} 💿 C. Breitenmoser-Würsten ¹⁰ 💿
L. Bufka ⁹ 💿 R. Černe ¹¹ 💿 J. Červený ⁸ 💿 N. Drouet-Hoguet ¹² 💿 M. Ďuľa ¹³ 💿
C. Fuxjäger ¹⁴ M. Herdtfelder ¹⁵ L. Hočevar ¹⁶ W. Jędrzejewski ^{17,18} R. Kont ¹⁹
P. Koubek ^{8,20} R. Kowalczyk ¹⁸ M. Krofel ¹⁶ J. Krojerová-Prokešová ^{20,21}
J. Kubala ^{22,23} D J. Kusak ²⁴ D M. Kutal ^{13,25} D J. D. C. Linnell ^{26,27} D J. Mattisson ²⁸ D
T. L. Middelhoff ⁵ D. Melovski ²⁹ 🖻 A. Molinari-Jobin ^{10,30} 🖻 J. Odden ³¹ 🖻
H. Okarma ³² 🖻 A. Ornicāns ⁷ 🖻 N. Pagon ¹¹ 🖻 J. Persson ⁶ 🗈 K. Schmidt ¹⁸ 🗈
M. Sindičić ²⁴ 💿 🕴 V. Slijepčević ³³ 👘 B. Tám ³⁴ 🕴 F. Zimmermann ^{10,35} 💿 👘
S Kramer-Schadt ^{3,36} M Heurich ^{1,2,27}

Correspondence

Marco Heurich, Wildlife Ecology and Conservation Biology, Chair of Wildlife Ecology and Management, Albert-Ludwigs-Universität Freiburg, Tennenbacher Straße 4, D-79106 Freiburg, Germany. Email: marco.heurich@wildlife.uni-freiburg.de

Article impact statement: Survival of Eurasian lynx across Europe is driven mainly by human-caused mortalities, in particular illegal killing.

Funding information

European Union INTERREG IV; MAVA Foundation: Norges Forskningsråd, Grant/Award Numbers: 251112, 281092; SCOPES programme; European Commission, Grant/Award Number: LIFE16 NAT/SL/000634 project 'LIFE Lynx'; Nature Protection Division of the County Governor's Office for Innlandet, Viken, Vestfold & Telemark, Trøndelag, Nordland, Troms & Finnmark Counties; Czech Academy of Sciences Institutional Research Plan; Swiss Foundation for the Scientific Research; Norwegian Directorate for Nature Management; Prof. Klaus Riepe und Ellen Riepe-Brunnström-Stiftung; Stotzer-Kästli Foundation; Deutsche Bundesstiftung Umwelt, Grant/Award Number: AZ 20017/504; People's Trust for Endangered Species; Agassiz Foundation; Nature Conservation Agency of the Czech Republic; Society for the Protection of Animals of the Canton

Abstract

Survival and cause-specific mortality rates are vital for evidence-based population forecasting and conservation, particularly for large carnivores, whose populations are often vulnerable to human-caused mortalities. It is therefore important to know the relationship between anthropogenic and natural mortality causes to evaluate whether they are additive or compensatory. Further, the relation between survival and environmental covariates could reveal whether specific landscape characteristics influence demographic performance. We used telemetry data on 681 Eurasian lynx (Lynx lynx), a model apex predator with large spatial requirements, that were tracked across their European distribution. Through time-to-event analyses, we sought to determine the variables associated with differences in their survival. Illegal killing was the main cause of mortality (33.8%), and mortality rates were similar in protected and hunted populations (8.6% and 7.0% per year, respectively). Survival varied greatly across populations (70-95% per year). Across all study sites, higher hunting and anthropogenic mortality rates were partially compensated by lower rates of other mortality causes but not by natural mortality alone. Variation in survival depended on sex (female survival was 1.5 times greater than male survival) and seasonality (highest risk during hunting season and winter), and lower survival rates were correlated with higher human modification of landscapes at both coarse (home range composition) and fine (habitat use within home range) scales. Some variation in survival was driven by unobserved factors, which, given the high rates of human-caused mortalities, including illegal killing, are of foremost concern. Due to the low natural mortality rates in protected and hunted populations, we conclude that anthropogenic causes of mortality are likely close to additive, such that maintaining or increasing refuge habitat with little human disturbance is critical to lynx conservation.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2025 The Author(s). Conservation Biology published by Wiley Periodicals LLC on behalf of Society for Conservation Biology.

and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

5231739, 0, Downloaded from https://conbio.onlinelibrary.wiley.com/doi/10.1111/cobi.14439 by Instytut Ochrony Przyrodyon, Wiley Online Library on [15/0]/2025]. See the Terms and Conditions (https://onlinelibrary.wiley.com/term)

of Vaud; Javna Agencija za Raziskovalno Dejavnost RS, Grant/Award Numbers: N1-0163, P4-0059; Polish Ministry of Science and Higher Education; Naturvårdsverket, Grant/Award Numbers: F-56-09, V-220-08; Operational Programme Environment through State Environmental Fund of the Czech Republic; European Union Interreg IIIA; WWF Switzerland; Hunting Administration of the Canton of Vaud, Fribourg and Bern; Swiss Federal Office for the Environment; Swiss League for Protection of Nature (Pro Natura); Svenska Forskningsrådet Formas, Grant/Award Numbers: 2010-1007, 2015-01207; Dr. Berthold Suhner Foundation; Operational Programme Research, Development and Education; Regina-Bauer Stiftung

INTRODUCTION

Survival is a key demographic rate, determined by underlying spatiotemporal distributions of risks and resources (Gaillard et al., 2010). Natural and anthropogenic mortalities vary by period and habitat, which increase or decrease survival probability (DeCesare et al., 2014). This means the survival of an individual should be correlated with the landscapes that they use (Fahrig, 2007). Connecting wildlife mortality events to landscape characteristics can, therefore, potentially reveal how species respond to different pressures (Bastianelli et al., 2021; Oriol-Cotterill et al., 2015). Conservation of threat-ened species hinges on understanding the factors, including anthropogenic, that influence survival and causes of mortality (Goodrich et al., 2008). These data are also necessary for evidence-based engagement with stakeholders in the context of human–wildlife coexistence (Redpath et al., 2013).

Wildlife survival rates in anthropogenic landscapes often depend on direct human actions. Behavioral plasticity can help offset anthropogenic disturbances, for example, by avoiding novel risk factors (Sih, 2013), but large carnivores must confront risks to maintain extensive home ranges that meet their energetic requirements (Carbone et al., 2005). In most cases, home ranges are fully outside protected areas, which could make avoidance of human-related risks impossible. Moreover, mortality risk can even be high in protected areas (Rauset et al., 2016). As a result, populations can exhibit sourcesink dynamics between areas of differing human influences. For example, occurrence of brown bear (Ursus arctos) in more human-influenced areas relies on immigration from less disturbed wilderness areas (Lamb et al., 2020), and pumas (Puma concolor) occupying remote areas are a source in more urbanized areas (Nisi et al., 2023).

Habitat selection is a hierarchical process that occurs at different scales to maximize fitness (Johnson, 1980). It is hypothesized that species will avoid the most limiting factors at coarser spatial scales, such as the selection of habitats for their home range composition (hereafter landscape) (Rettie & Messier, 2000). At finer selection scales, such as use of habitat within the home range (hereafter home range), individuals may control the remaining risks spatially and temporally and focus on other basic needs, such as prey resources (Filla et al., 2017; Suraci et al., 2019). However, whether this hierarchy is also reflected in individual fitness is not guaranteed (Basille et al.,

KEYWORDS

cause-specific mortality, compensatory mortality, Eurasian lynx, large carnivore, Lynx lynx, survival

2013). To understand the drivers of survival and mortality, they must be studied across gradients of risks and resources and at different spatial scales.

A key concept in wildlife management is that some proportion of individuals in a natural population are destined to die naturally (Errington, 1956). This gives rise to the classic hypothesis that if increases in a given mortality source can be compensated by decreases in other mortality sources, then survival remains constant (i.e., compensatory mortality) (Burham & Anderson, 1984). However, mortality causes can also be additive, whereby increases of a given mortality cause are directly proportional to decreases in survival, or partially compensatory, whereby reductions in other mortality causes do not fully compensate and survival decreases (e.g., Sandercock et al., 2011). Characterizing the relationship between survival and different mortality causes poses a fundamental question in population ecology (Boyce et al., 1999) and is important to ensure sustainable harvest of game species (Marboutin et al., 2003; Medellín, 1999). Due to their ecology, large carnivores are particularly vulnerable to human-caused mortalities (Krofel et al., 2015), which are often more frequent than natural causes (Moss et al., 2016). In hunted and protected populations, other anthropogenic mortality causes may have additive or compensatory relations to natural mortality (e.g., Murray et al., 2010). By exploring how survival is related to different mortality factors, one can better understand species' vulnerability to demographic pressures and provide important information for conservation of protected populations and sustainability in hunted populations (e.g., Wolfe et al., 2015).

Assessment of large carnivore survival has typically been conducted on a local scale. Such studies of single populations (e.g., Suutarinen & Kojola, 2017) provide locally relevant information, but their application to interregional questions may be limited. Studies of survival and mortality in multiple populations, or study areas, covering gradients of environmental conditions and management regimes (e.g., Smith et al., 2010; Wolfe et al., 2015) can improve understanding of where large carnivore populations might struggle to coexist with humans (Gilroy et al., 2015). Combining data from different regions can also help improve understanding of the interplay between survival rates and different mortality causes at a species distribution level. These outputs could inform conservation plans at geographic scales that are meaningful for species whose populations cover vast extents, such as large carnivores (Benson et al., 2023).

Telemetry tracking is a precise way to estimate survival in wild animal populations because individuals' fates are known (Murray & Patterson, 2006) and all mortality causes have equal detection probabilities (Naef-Daenzer et al., 2017). Analyses of tracking data represent a special subclass in survival modeling, formally called known-fate models (Heisey & Fuller, 1985), as opposed to methods where mortality must be partially inferred (e.g., camera trapping) (Saracco et al., 2010). Known-fate models can provide statistically robust estimates of survival and mortality because they consider the number of animals tracked and the length of time they were tracked (Heisey & Fuller, 1985). Multivariate known-fate modeling has, therefore, frequently been used to relate observed covariates, such as sex and landscape composition, from telemetry locations to survival (Basille et al., 2013; Brodie et al., 2013). The heterogeneity of survival not captured by spatial variation in observed covariates (e.g., effects of human acceptance, wildlife population density) may also contain important information (Halstead et al., 2012). Survival models that can delineate the effects of both observed and unobserved spatially dependent factors (e.g., Zhou & Hanson, 2015) would be advantageous for assessing wildlife survival across large geographic extents because unexplained variability is considered. Thereby, the common drivers, which can inform decision-making, and variability of unknowns can be better understood.

We compiled telemetry data on 681 Eurasian lynx (Lynx lynx) (hereafter lynx), a model apex predator with large spatial requirements, that were tracked across their European distribution (Figure 1a). We used time-to-event analyses to provide survival and cumulative incidence of cause-specific mortality rates for lynx across Europe (Figure 1b). These quantities are vital for evidence-based population viability analyses. Further, we used these to test for consistency with competing mortality hypotheses, hypothesizing, in line with the compensatory hypothesis, that higher anthropogenic (H1) and hunting mortality (H2) are not associated with lower overall survival and therefore natural mortality declines as anthropogenic (H3) and hunting mortality (H4) declines, and higher hunting mortality is associated with lower alternative mortality causes (H5) and less illegal killing (H6). Further, we aimed to improve species-level understanding of the factors driving survival with multivariate models (Figure 1c) to inform management as to how to spatially allocate conservation resources more effectively. According to the limiting factor hypothesis in hierarchical habitat selection (Rettie & Messier, 2000), we hypothesized that selected landscape components affect survival in a scale-dependent manner (H7). At the coarser landscape scale (i.e., home range composition), we hypothesized that human habitat modification and disturbances have a strong negative effect on lynx survival (H8). At the finer home range scale (i.e., within home range habitat use), we hypothesized that survival is less correlated to human-related landscape characteristics (H9) because these are already avoided at the coarse scale and survival is more positively affected by landscape characteristics indicative of prey resources (H10). Finally, we hypothesized that high-quality habitat has a positive effect on survival at both spatial scales (H11).

METHODS

Data acquisition

We collected tracking data from 21 telemetry studies, representing 9 out of 10 populations occurring in Europe. A total of 681 lynx individuals (314 females, 367 males) were tracked for a combined 1052 years (Appendix S1; Figure 1a). Animal captures and experimental procedures were approved by the government agencies and ethics committees relevant to each study area in accordance with their respective animal welfare laws (details in Appendix S2).

At first capture, 327 (48%) individuals were adults (>2 years), and 115 (17%) and 239 (35%) were subadults (1-2 years) and juveniles (<1 year), respectively. We assigned age classes with high confidence due to the synchronized nature of birth dates (Mattisson et al., 2022), body size development, and social-spatial behaviors (i.e., mating, dispersal). We updated individuals' age classes over time based on their age, which we calculated from the observed, or estimated, birth date. We set the estimated birth dates as the start date of the birthing season (1 May) (Mattisson et al., 2020) and calculated the year via the individual's age class at the time of their first observation (e.g., Weingarth et al., 2012) or postmortem dental analysis (e.g., Marti & Ryser-Degiorgis, 2018). Birth years varied from high confidence (natal den visits, camera trapping of females with kittens, postmortem) (54%) to medium confidence (juvenile or subadult at first observation) (23%) to low confidence (adults at first observation, i.e., minimum age) (23%). We conducted all analyses with respect to age class to ensure robust results, and some aspects were repeated with age (see time scales below). We acquired lynx location data, by triangulation of radio signals for very high frequency (VHF) collars, and remote or direct download for global positioning system (GPS) collars, with an average of 1.6 and 3.8 locations/day, respectively. Overall, our data set included 219 mortality events, and the oldest lynxes, based on telemetry tracking observations, were 18.0 and 14.7 years for females and males, respectively.

We investigated potential mortality following triggering of on-collar mortality sensors or immobility. We categorized telemetry tracking that ended due to mortality into the following causes: natural (disease, inter- or intraspecific killing, starvation, falls, and unknown natural causes), illegal killing, legal killing, vehicle collisions (road and train), and unknown (e.g., carcass decomposed or consumed). We included suspected illegal killing following Andrén et al.'s (2006) criteria to distinguish illegal killing from radio transmitter failure and unknown fates (Appendix S3). We also considered the broad groups of anthropogenic (illegal killing, legal killing, and vehicle collisions) and nonhunting (natural, illegal killing, vehicle collisions, and unknown) mortalities for hypotheses H1, H3, and H5. We rightcensored telemetry tracking that ended without mortality (i.e., survival time was not observed because the tracking ended before the animal died; collar was removed, dropped off, or failed).



FIGURE 1 Workflow in a study of Eurasian lynx survival in Europe: (a) acquisition of lynx tracking data and geospatial data (¹, Table ; ², Appendices S7 & S8; ³, Figure 2; ⁴, Figure 3; ⁵, Appendix S5; ⁶, Appendix S1; ⁷, Table 2; ⁸, Table 3 and Appendix S10; ^d, part [d] of this figure; ^e, part [e] of this figure; GPS, global positioning system; VHF, very high frequency), (b) tracking period observations (i.e., start time, end time, fate) to provide point estimates and curves of survival and cumulative incidence of cause-specific mortalities based on product-limit estimation (i.e., KM, Kaplan–Meier) used also in the analysis of mortality hypotheses, (c) combining tracking periods with location data to model the effects of landscape covariates on survival rates (HR, home range; IID, independent and identically distributed; DIC, deviance information criterion; LPML, log pseudo-marginal likelihood; GRF, Gaussian random field), (d) transformation of tracking periods to different time scales (d.1, age time scale; d.2, annual time scale), and (e) procedure used to connect season (age time scale), hunting periods (annual time scale), and landscape covariate; val., value).

We transformed the dates and times of the data to comply with 2 different survival time scales (Figure 1d). First, we created a continuous age time scale, which is the natural scale for ageing with the time origin at birth (Lamarca et al., 1998), whereby observations are intervals of age conditional on survival from birth until the first capture, known as left truncation or delayed entry (Geskus, 2011). Second, we created a recurrent annual time scale with origin 1 May (Fieberg & DelGiudice, 2009). The annual time scale splits tracking periods on 1 May so that they represent interval times with a possible range of 0–365 days, regardless of the year. This is useful for estimating annual rates for management types and age classes. We conducted statistical analyses with R 4.1.3 (R Core Team, 2022) and visualized results with the package ggplot2 (Wickham, 2016).

Nonparametric survival and cause-specific mortality estimates

Nonparametric methods are appropriate to estimate empirical point estimates and curves of survival rates and cumulative incidence rates of cause-specific mortality with few assumptions (Figure 1b). We estimated these quantities for sex, management type (legally hunted or protected), and study area. We estimated the survival rates for each year of age based on the age time scale and estimated annual means for subadult and adult age classes based on the annual time scale. We used the product-limit (i.e., Kaplan-Meier) estimator (Kaplan & Meier, 1958) from the package survival (Therneau, 2021) to estimate survival rates. We used weighted product-limit estimation to estimate cumulative incidence rates of competing mortalities while accounting for left truncation (Geskus, 2011). For this, with the package mstate (de Wreede et al., 2011), we replicated the data set for each mortality cause and augmented it with time-dependent truncation weights and censoring weights that accounted for the competing events. We calculated conditional survival rates with the package condsurv (Zabor & Ganon, 2019) and visualized curves with survminer (Kassambara et al., 2017).

Additive versus compensatory mortality

To assess our predictions regarding the mortality hypotheses (H1-H6), we used nonparametric estimates of annual survival and mortalities for adults only from the study areas where survival rates could be estimated (i.e., survival <1). We fitted a regression model (response vs. predictor) for each hypothesis: H1, survival versus anthropogenic factors (hunting, illegal killing, vehicle collisions); H2, survival versus hunting; H3, natural mortality versus anthropogenic mortality; H4, natural mortality versus hunting mortality; H5, nonhunting mortality versus hunting mortality; and H6, illegal killing versus hunting mortality. For H1 and H2, we predicted regressions with no trend (fully compensatory) (Appendix S4), whereas for H3-H6, we predicted negative trends, which would show a decline in the response mortality rate as their respective predictor mortality increases (compensation). We used beta regression models from betareg (Cribari-Neto & Zeileis, 2010) because they are

suitable for modeling responses in the range 0–1. To account for the uncertainty in the nonparametric estimates, we sampled 10,000 values for each rate from beta distributions (mean and variance defined by each rate's estimate and standard error squared, respectively) and fitted a regression for each sample. Thereafter, we estimated the mean β and 90% highest posterior density intervals (HPD) across samples (Benson et al., 2023) to evaluate H1–H6. For H1 and H2, we assessed the trends and intercepts of the 10,000 regression lines for consistency with the hypotheses: compensatory and overcompensatory (positive trend), additive and superadditive (negative trend with slope $\langle -S_0 \rangle$, and partially compensatory (negative trend but with slope $\rangle -S_0$), for which we assumed the baseline survival rate (no hunting), S_0 , was the intercept of each regression (Appendix S4).

Multivariate survival modeling

Multivariate models are an appropriate way to make statistical inference about individuals' mortality risk depending on their exposure to multiple time-varying and static factors and treatments (e.g., Bradburn et al., 2003). These factors are quantified, or measured, when an individual is observed at certain instances during their monitoring. Between these instances, it is normally assumed that these factors do not change in value and that the risk associated with them during this discrete period is constant (i.e., piecewise constant hazards). In this way, we related observations of lynx's survival and mortality with temporal exposure to landscape features and seasons during their tracking (Figure 1e), as well as their sex. Season is expected to affect survival of lynx (e.g., Andrén et al., 2022); therefore, to accommodate the piecewise constant hazards assumption, we split tracking periods by season (spring, March-May; summer, June-August; autumn, September-November; winter, December-February) and by country-specific hunting periods (Figure 1e; Appendix S5). Spring coincides with the onsets of the birthing period and natal dispersal of subadults, summer with weaning of kittens, autumn with higher female mobility with kittens, and winter with the onset of mating (Breitenmoser-Würsten et al., 2007; Mattisson et al., 2022; Samelius et al., 2012; Zimmermann et al., 2005). We fitted candidate models at each spatial scale and survival time scale, divided into 3 sets: base, habitat suitability (HSI), and component models, with different covariates described below.

Landscape usage and covariates

We used individuals' locations collected during each season's tracking to characterize their landscape use (i.e., the areas or locations individuals were exposed to over time). These are corollaries to the scales in hierarchical habitat selection (Johnson, 1980), specifically: landscape scale and home range scale (e.g., Oeser, Heurich, Kramer-Schadt, Mattisson, et al., 2023. For the home range scale, we assigned each individual's locations to seasonal tracking periods by their acquisition times. For the landscape scale, we took all the locations assigned

within each individual's seasonal tracking periods and performed kernel density estimation (KDE), implemented in the package amt (Signer et al., 2019); we used the 95% vertices. Similar to Andrén et al. (2022), for periods when individuals had fewer than 25 locations (54%), we used sex-specific average daily distance traveled as a radius around locations to delineate landscape scale use (Figure 1e). We used different radii for 3 latitudinal groups: above 65°N (female, 4.4 km; male, 6.9 km), from 55 to 65°N (female, 3.2 km; male, 4.6 km), and below <55°N (female, 2.1 km; male, 3.5 km). If an individual had zero locations in a given period, we brought the previous period forward (10% of cases).

We compiled covariates to characterize landscape composition. For the HSI models, we used the HSI map derived in a Europe-wide study by Oeser, Heurich, Kramer-Schadt, Mattisson, et al. (2023). For the component model set, we collected variables that corresponded to habitat or anthropogenic influences: forest integrity (index of human modification of forests), greenness variability (vegetation seasonality), topographic ruggedness, land cover (individually as proportions of forest, shrub, grass, crop, and urban land covers), human modification index, accessibility (travel time to cities), distance to major roads, distance to minor roads, distance to settlements, and human population density (Appendix S5). We extracted covariate values at both spatial scales, taking the mean of each for every individual's seasonal tracking period, thereby landscape covariates varied temporally. To avoid potentially problematic correlations between covariates and reduce the dimensionality (component model set), we used principal component analyses (PCAs) and retained principal components (PCs) that explained at least 5% of the variance (Appendix S6) and fitted a model for each, as well as one model with the first 2 PCs that explained the most variance (landscape, 50.3% and 17.5%, respectively; home range, 44.2% and 12.5%, respectively).

To these, we added categorical covariates (base model) parametrically, including sex (time invariant; 2 levels [female as reference and male]) for all models and additional time-scalespecific covariates. For the age time scale models, we included season (time varying; 5 levels, autumn, winter, hunting, spring, summer) due to annually varying risk (Fieberg & DelGiudice, 2009). We chose autumn as the reference level due to demographic stability typical in this season (e.g., Weingarth et al., 2015). We assigned hunting season to individuals exposed to this risk by splitting tracking periods at the start and end of the country-specific hunting periods (Appendix S1) and replacing the calendar season within this period with the level of hunting (Figure 1e). For the annual time scale models, we included age class (time varying; 3 levels, juvenile [reference], subadult, adult) and hunting period (time varying; 2 levels, hunting [reference], nonhunting), whereby we assigned the level nonhunting to all tracking periods outside hunting periods.

Model fitting and comparison

To determine the effects of time-varying covariates at different spatial scales on survival (H7–H11), we fitted multivariate

Bayesian semiparametric accelerated failure time models (Zhou & Hanson, 2018) from the package spbayessurv (Zhou et al., 2020). This is a powerful method that accommodates arbitrary censoring, left truncation, time-varying covariates, and geographic information without the constraint of proportional hazards. In the absence of precise information, we used noninformative default priors. We fitted models with adaptive Markov chain Monte Carlo sampling methods, comprising an initial parametric phase with a lognormal centering distribution (5000 draws) that provides a guide to the baseline survival function in the main semiparametric phase (5000 burn-ins, retaining 10,000 draws from 50,000). We included frailty terms, which are equivalent to random effects, to account for unobserved factors that influenced survival and to address the lack of independence of individuals with repeated observations (annual time scale) and multiple individuals in the same study area (age and annual time scales). Specifically, we used independent and identically distributed (IID) frailties for nonspatially referenced models, where observations are grouped by individual and study area, and Gaussian random field (GRF) frailties for spatially explicit models, where variation of risk depends on a multivariate spatial distribution. We used the centroids of individuals' locations as the input coordinates for the GRF frailty.

We fitted each model set once for each frailty (GRF, IID), survival time scale (age, annual), and spatial scale (landscape, home range) for a total of 64 models. We used the deviance information criterion (DIC) and the log pseudo-marginal likelihood (LPML), where smaller DIC and larger LPML indicate better model performance, to select the most parsimonious models (Δ DIC or Δ LPML <2) in each time and spatial scale combination (16 candidate models each). We checked model performance of selected models visually via conditional Cox–Snell residuals (Zhou & Hanson, 2018). To determine coefficient effects, we estimated their evidence ratios relative to zero for the continuous covariates and among different levels of the categorical covariates with the package brms (Bürkner, 2017).

RESULTS

Survival estimates and cause-specific mortality estimates

Overall, the median survival age (i.e., age when survival probability reaches 0.5) of males was 2.59 (95% confidence interval [CI] 1.43–3.43) and 2.68 (95% CI 1.39–6.33) years in hunted and protected populations, respectively, and for females, it was 4.00 (95% CI 3.28–5.85) and 3.36 (95% CI 2.15–11.39) years, respectively (Figure 2a). The lifespans of females and males in protected populations were generally longer than their hunted counterparts, though annual adult survival had a large range from 0.70 (Harz Mountains) to 0.95 (Dinaric southeastern Alps). In summary, the point estimates showed that hunting and illegal killing were the most important mortality causes for lynx and the incidence of illegal killing in protected and hunted populations was almost equal to the incidence of hunting mortality in

(a) Female hunted (FH) Female protected (FP) Male hunted (MH) ---- Male protected (MP) 1 00 0.75 Survival probability 0.50 0.25 н "MP FH мн 'FP 0.00 10 7 9 11 12 6 8 13 14 15 Age (years) (b) (c) 0.8 IK Cumulative incidence IK 0.6 Ν н Ν 0.4 U υ 0.2 VC VC 0.0 10 15 5 10 15 0 Age (years) Age (years) Hunting (H) Illegal killing (IK) Natural (N) Unknown (U) — Vehicle collision (VC)

FIGURE 2 For Eurasian lynx in Europe, (a) product-limit (Kaplan–Meier) survival estimates for sex and management groups (dotted lines, median survival ages) and weighted product-limit (Geskus, 2011) cumulative incidence estimates of cause-specific mortalities of (b) lynx subject to hunting and (c) protected lynx (no legal hunting besides extenuating circumstances). Tabularized values are in Appendices \$7 and \$8.

hunted populations (point estimates, Table 1; curves, Figure 2; detailed groups, Appendices S7 & S8).

Additive versus compensatory mortality

In contrast to our hypotheses (H1 & H2), there was no evidence of compensatory mortality, as annual adult survival decreased with increasing anthropogenic mortality ($\beta = -3.67, 90\%$ HPD -6.23 to -1.22) and hunting mortality ($\beta = -4.08, 90\%$ HPD -5.90 to -2.01) (Figure 3a,b). Instead, these results support the additive or partially compensatory mortality hypotheses because survival was negatively influenced by anthropogenic and hunting mortality. Based on the 10,000 sampled regressions, 22% and 46% of the regression slopes for anthropogenic and hunting mortality against survival, respectively, were steep enough to fit the additive mortality hypothesis. The remaining majority were consistent with partially compensatory mortality (Appendix S9). We found negative relationships between natural mortality and anthropogenic ($\beta = -2.16, 90\%$ HPD -6.11

to 2.22) and hunting ($\beta = -2.44$, 90% HPD -5.46 to 0.946) mortality rates (Figure 3c,d). However, their HPD intervals overlapped zero and, therefore, provided no evidence for the compensatory mortality we predicted (H3 & H4). This suggests that anthropogenic and hunting mortalities were not, or only partially, compensated by changes in natural mortality. Indeed, adult natural mortality rates were similar in hunted and protected populations (Table 1). The posterior mean regression of nonhunting mortality with hunting mortality showed a negative trend ($\beta = -0.944$, 90% HPD -2.87 to 1.21), consistent with H5. There was a slight positive trend in the regression between mortality due to illegal killing and hunting mortality $(\beta = 1.11, 90\% \text{ HPD} - 1.72 \text{ to } 5.25)$, in opposition to H6. However, in both cases, the regressions were statistically inconclusive because their HPD intervals overlapped zero (Figure 3e,f). In summary, increases in hunting mortality were not directly proportional to reductions of any single mortality cause or the combination of all nonhunting mortalities across the different study areas. This suggests that populations were only partially able to compensate for hunting mortality.

Conservation Biology



FIGURE 3 The relationships between Eurasian lynx (a, b) survival and (c, d) natural mortality with hunting mortality and anthropogenic mortality, respectively, and the relationship between (e) nonhunting mortality and (f) illegal killing, with hunting mortality used to test the hypotheses of compensatory mortality (H1–H6) (black dots, nonparametric point estimates; vertical and horizontal lines, 95% confidence interval of adult annual survival and annual cumulative incidence mortality rates, respectively; red lines, mean regressions; pink, 90% highest posterior density intervals [HPD] based on 10,000 samples [blue dots]; gray lines, regression of each sample).

TABLE 1 Annual nonparametric estimates of Eurasian lynx survival and cumulative incidence of competing cause-specific mortality events for a subset of groups.

D		0	Management or		
Parameter	Age class	Sex	other	Estimate (95% CI)	
Survival	Adult	Female	Hunted	0.82 (0.78–0.86)	
			Protected	0.86 (0.79–0.91)	
		Male	Hunted	0.76 (0.70-0.80)	
			Protected	0.83 (0.76–0.88)	
	Subadult	Female	Hunted	0.93 (0.83–0.97)	
			Protected	0.72 (0.50-0.85)	
		Male	Hunted	0.73 (0.58–0.84)	
			Protected	0.78 (0.55-0.90)	
Natural	Adult	Female	Protected	0.031 (0.0010-0.060)	
		Male	Protected	0.043 (0.0090-0.077)	
		Mean	Hunted	0.033 (0.017–0.048)	
			Protected	0.038 (0.014-0.061)	
	Subadult	Mean	Hunted	0.037 (0.0010-0.071)	
			Protected	0.111 (0.014–0.19)	
Illegal killing	Adult	Female	Protected	0.081 (0.031-0.13)	
		Male	Protected	0.089 (0.041–0.13)	
		Mean	All populations	0.074 (0.056–0.091)	
			Hunted	0.070 (0.048-0.091)	
			Protected	0.086 (0.051-0.12)	
			Reintroduced	0.079 (0.041-0.12)	
	Subadult	Mean	Hunted	0.050 (0.0060-0.092)	
			Protected	0.098 (0.0020-0.186)	
Hunting	Adult	Female	Hunted	0.055 (0.028-0.080)	
		Male	Hunted	0.15 (0.10-0.20)	
Vehicle		Mean	Hunted	0.012 (0.002–0.022)	
			Protected	0.012 (0-0.025)	
	Subadult	Mean	Hunted	0.022 (0-0.051)	
			Protected	0.048 (0-0.11)	

Note: For nonparametric estimates calculated for groups not shown, see Appendices S7 and S8.

Abbreviation: CI, confidence interval.

Covariate effects

Across all selected models, we found very strong evidence that male survival times were lower than female survival times (30– 36% less) (Table 3). Naturally, survival declined as lynx aged; the highest hazard rates were from 1 to 3 years (Appendices S12 & S13). Considering seasonality, there was very strong evidence across models that hunting season reduced survival probabilities more than any calendar season. The age time scale models offered strong evidence that survival was lower during winter compared with autumn and summer but only moderate evidence compared to spring. Based on the age time scale models, lynx survival probability declined 3 times faster during hunting season (typically winter) and 1.7 times faster during winter Conservation Biology 🔇

(without hunting) than during autumn. There was weak to no evidence that spring, summer, and autumn differed. The annual time scale models reflected these results with hazard rates generally increasing in winter (Appendices S14 & S15). Considering the selected models fitted with the annual time scale, there was only a little evidence (ratios 4–6) that juvenile, subadult, and adult lynx differed in annual survival probabilities. This is likely due to the smaller sample size of younger lynx and because the annual time scale does not consider senescence, which can cause bias, especially for the adult age class (Koons et al., 2014), whereas the alternative age time scale showed senescence naturally.

At the landscape scale, several models with landscape covariates were selected (Table 2), but most found only little to no evidence that they affected survival rates (Table 3). There was moderate evidence from one model on the annual time scale that PC1, which primarily described a gradient from human-dominated areas to areas with low accessibility and great distance from human infrastructure (Appendix S16), was associated with higher survival times. Higher values of PC1, and therefore home range composition with little anthropogenic influence on average, increased survival probability (H8). Although one HSI model was selected, there was little evidence that the variable affected survival at this spatial scale (H11).

At the home range scale, models including landscape covariates were also selected (Table 2). There was moderate to strong evidence on the age time scale and strong evidence on the annual time scale that the landscape variability described by PC1 affected survival. The PC1 characterizes a gradient from highhuman-modification areas, including high forest land cover, to areas with more shrub and grass land covers farther from human settlements and roads (Appendix S17; H9 & H10). One model selected on the annual time scale provided weak to moderate evidence that if lynx used locations with on average higher habitat suitability, their survival probability would be higher (H11). Finally, one selected home range model showed weak evidence that lower human populations and urban cover could increase survival times. The evidence that landscape characteristics affected survival was stronger at the home range scale (i.e., higher posterior probabilities of estimates in the selected models, H7).

Considering the selected models that included frailty as a spatially correlated risk (GRF frailty), we found the effects posed by unobserved drivers of survival (i.e., those not explained by the covariates) were relatively restricted spatially (Table 3). This means that the combined effects of unexplained factors that caused variation in survival were similar for individuals found close together (e.g., approximately 5% lower correlation at 10 km). The similarity of these effects decreased rapidly with increasing distance between individuals (e.g., approximately 50% lower correlation at 150 km). There was also a relatively high variance in this component, indicating considerable heterogeneity in the unobserved factors (τ^2 from 0.1 to 0.9).

The selected models were consistent with statistical modeling framework assumptions (Appendices \$18-\$21), and the

Conservation Biology

Comp. GRF, PC1

Comp. GRF, PC1

Comp. GRF, PC4

Comp. GRF, PC2

Comp. GRF, PC1

Comp. GRF, PC1 + 2

Base GRF

Base IID

Base GRF

Base IID

HSI GRF

Base GRF

Base IID

1.26

1 27

5.99

0.24

0.96

0.00

1.15

5.85

0.00

1.19

1.27

3.69

6.71

Т sc

Home range, age

Home range, annual

TABLE 2 Comparisons of multivariate models of survival of Eurasian lynx for selected ^a and base ^b candidate models for each spatial scale and survival time scale ^c .						
Scale	Model	DIC ^d	ΔDIC	LPML ^e	ΔLPMI	
Landscape, age	Comp. GRF, PC5	6438.70	0.00	-3229.46	1.96	
	Comp. GRF, PC1	6439.16	0.46	-3227.50	0.00	
	Comp. GRF, PC4	6439.70	0.99	-3228.75	1.25	
	HSI GRF	6439.84	1.13	-3229.36	1.86	
	Base GRF	6444.21	5.51	-3228.37	0.87	
	Base IID	6450.44	11.74	-3232.33	4.83	
Landscape, annual	Comp. GRF, PC5	3455.60	0.00	-1733.87	4.00	
	Comp. GRF, PC4	3455.93	0.33	-1729.87	0.00	

3457.24

3457.94

3465.11

6434.54

6436.71

6436.98

6441.09

6446.83

3452.99

3453.24

3454.92

3461.04

3465.16

1.64

2.33

9.51

0.00

2.16

2.44

6.55

12.28

0.00

0.25

1.93

8.05

12.17

Abbreviations: Comp., component; GRF, Gaussian random field frailties; HSI, habitat suitability index; IID, independent and identically distributed frailities; PC, principal component.
Models selected with change in deviance information criterion (Δ DIC) or change in log pseudo-marginal likelihood (Δ LPML) <2. Comparison of all candidate models is in Appendix S1
Base covariates only: sex + season (age time scale) and sex + hunting + age class (annual time scale).

^cLandscape and home range spatial scales and age and annual survival time scales.

^dDeviance information criterion. The smaller the DIC, the better the model quality. Models ordered by increasing DIC.

^eLog pseudo-marginal likelihood. The larger the LPML, the better the model performance.

coefficient estimates were stable for all covariates with weak to very strong evidence (Table 3; Appendix S11).

DISCUSSION

Making use of a large telemetry data set, we provided the first continental-scale assessment of survival rates and cumulative incidence of cause-specific mortalities for a large carnivore in the human-dominated landscapes of Europe. This offers managers and researchers the most up-to-date knowledge on lynx survival probability. We found that human-related mortalities, including illegal killing, legal hunting, and vehicle collisions, exceeded natural mortality causes, as was previously shown in some studies focused on specific lynx populations (Andrén et al., 2006; Breitenmoser-Würsten et al., 2007; Schmidt-Posthaus et al., 2002; Sindičić et al., 2016). Illegal killing was the most important source of mortality risk for lynx in protected populations. In hunted populations, illegal killing posed a similar risk as legal hunting, and lynx were similarly likely to die naturally. Indeed, across our study sites, we found strong evidence

that anthropogenic and hunting mortalities were only partially compensated by reductions in natural mortality or other causes. Based on multivariate survival models, we found lynx had an approximately 20% probability of reaching 15 years of age. Survival was mostly driven by sex, season, and landscape characteristics. For both landscape and home range scales, lynx that used areas with fewer anthropogenic influences farther from human infrastructure were likely to survive longer, provided unobservable local conditions, such as illegal killing, were not detrimental.

-1731.12

-1731.14

-1735.86

-3227.12

-3227.84

-3226.88

-3228.04

-3232.73

-1729.59

-1730.79

-1730.87

-1733.28

-1736.30

As we showed for lynx, illegal killing is a prominent concern for many large carnivore species (Carter et al., 2017). In this context, sustainable harvest is often seen as a method to balance contrasting stakeholder wishes in the political arena (Linnell et al., 2010). In our study areas, hunting mortality was at best only partially compensated by lower mortality due to other causes. This was also the case when all anthropogenic mortality causes were pooled. Natural mortality occurred at similar rates in hunted populations and was not low enough to compensate fully. Indeed, lower rates of all other mortality causes were also not sufficient. Therefore, it is likely that natural mortality rates

Conservation Biology

 ${\bf TABLE \ 3} \quad {\rm Covariate \ effect \ estimates \ from \ selected^a \ multivariate \ Eurasian \ lynx \ survival \ models \ for \ each \ spatial \ and \ time \ scale \ with \ estimates \ from \ the \ most \ parsimonious \ model \ in \ which \ each \ covariate \ appeared^b. }$

Scale	Model	Term	Estimate ^c	90% CI ^d	Test ^e	Evidence ^f	₽ ^g
Landscape, age	Comp. GRF, PC5	eta sex: male	0.410	0.188 to 0.633	>female ^h	1249	0.999
		eta season: hunting	1.12	0.734 to 1.59	>autumn ^h	Inf	1.00
			0.645	0.236 to 1.12	>winter	302	0.996
			1.20	0.613 to 1.98	>spring	4999	0.999
		eta season: spring	0.320	-0.187 to 0.837	>autumn ^h	6.11	0.859
			0.178	-0.476 to 0.743	>summer	2.35	0.701
		eta season: summer	-0.0548	-0.470 to 0.377	<autumn<sup>h</autumn<sup>	1.48	0.598
		eta season: winter	0.647	0.243 to 1.11	>autumn ^h	399	0.997
			0.558	0.00911 to 1.19	>spring	19.2	0.950
			0.736	0.263 to 1.21	>summer	139	0.992
		β PC5	-0.00281	-0.209 to 0.203	<intercept< td=""><td>1.05</td><td>0.512</td></intercept<>	1.05	0.512
		arphi GRF frailty scale	0.00423	0.00149 to 0.00925			
		$ au^2$ frailty variance	0.378	0.143 to 0.759			
	Comp. GRF, PC1	β PC1	-0.0280	-0.125 to 0.0842	<intercept< td=""><td>2.17</td><td>0.684</td></intercept<>	2.17	0.684
	Comp. GRF, PC4	β PC4	0.0884	-0.0466 to 0.224	>intercept	6.34	0.863
	HSI GRF	β HSI	0.00491	-0.00380 to 0.0152	>intercept	3.88	0.795
Landscape, annual	Comp. GRF, PC5	eta age class: subadult	-0.359	-1.00 to 0.276	<juvenile<sup>h</juvenile<sup>	4.45	0.816
		eta age class: adult	-0.354	-1.00 to 0.240	<juvenile<sup>h</juvenile<sup>	4.89	0.830
			0.00494	-0.405 to 0.405	>subadult	1.06	0.515
		eta sex: male	0.337	0.0856 to 0.589	>female ^h	77.1	0.987
		$oldsymbol{eta}$ hunt. P ⁱ .: nonhunt.	-1.64	-4.33 to -0.632	<hunting<sup>h</hunting<sup>	Inf	1.00
		β PC5	-0.0168	-0.233 to 0.209	<intercept< td=""><td>1.28</td><td>0.562</td></intercept<>	1.28	0.562
		arphi GRF frailty scale	0.00306	0.000808 to 0.00648			
		$ au^2$ frailty variance	0.634	0.200 to 1.31			
	Comp. GRF, PC4	β PC4	0.125	-0.00313 to 0.261	>intercept	17.3	0.945
	Comp. GRF, PC1	β PC1	-0.0733	-0.142 to -0.00489	<intercept< td=""><td>25.5</td><td>0.962</td></intercept<>	25.5	0.962
	Base GRF	_					
Home range, age	Comp. GRF, PC1	eta sex: male	0.400	0.158 to 0.659	>female ^h	269	0.996
		eta season: hunting	1.10	0.710 to 1.56	>autumn ^h	Inf	1.00
			0.554	0.195 to 0.953	>winter	255	0.996
			0.375	-0.0385 to 0.845	>spring	12.8	0.927
		m eta season: spring	0.177	-0.283 to 0.624	>autumn ^h	2.82	0.738
			0.287	-0.188 to 0.727	>summer	5.25	0.840
		eta season: summer	-0.110	-0.527 to 0.293	<autumn<sup>h</autumn<sup>	2.06	0.674
		eta season: winter	0.553	0.141 to 1.01	>autumn ^h	80.9	0.987
			0.375	-0.0385 to 0.845	>spring	12.8	0.927
			0.663	0.255 to 1.09	>summer	587	0.998
		β PC1	-0.118	-0.224 to -0.0185	<intercept< td=""><td>49.7</td><td>0.980</td></intercept<>	49.7	0.980
		arphi GRF frailty scale	0.00323	0.000508 to 0.00788			
		$ au^2$ frailty variance	0.607	0.222 to 1.50			
	Comp. GRF, PC1 + 2	β PC2	0.00440	-0.0847 to 0.0875	>intercept	1.25	0.556
	Base GRF	-					
	Comp. GRF, PC4	β PC4	-0.0881	-0.182 to 0.0114	<intercept< td=""><td>13.0</td><td>0.928</td></intercept<>	13.0	0.928
Home range, annual	HSI GRF	eta age class: subadult	-0.360	-0.978 to 0.211	<juvenile<sup>h</juvenile<sup>	5.57	0.847
		eta age class: adult	-0.304	-0.828 to 0.213	<juvenile<sup>h</juvenile<sup>	4.78	0.827
			0.0555	-0.304 to 0.422	>subadult	1.44	0.591

TABLE 3 (Continued)

Scale	Model	Term	Estimate ^c	90% CI ^d	Test ^e	Evidence ^f	₽ ^g
		eta sex: male	0.340	0.110 to 0.587	>female ^h	105	0.990
		eta hunt. P.: nonhunt.	-1.01	-2.04 to -0.478	>hunting ^h	Inf	1.00
		β HSI	-0.00773	-0.0167 to 0.00108	<intercept< td=""><td>12.7</td><td>0.927</td></intercept<>	12.7	0.927
		arphi GRF frailty scale	0.00181	0.000598 to 0.00392			
		$ au^2$ frailty variance	0.846	0.282 to 1.75			
	Comp. GRF, PC2	β PC2	0.0597	-0.0323 to 0.159	>intercept	6.14	0.860
	Comp. GRF, PC1	β PC1	-0.122	-0.208 to -0.0515	<intercept< td=""><td>453</td><td>0.997</td></intercept<>	453	0.997

Abbreviations: Comp., component; GRF, Gaussian random field frailties; HSI, habitat suitability index; IID, independent and identically distributed frailties; PC, principal component. ^aCandidate models with change in deviance information criterion (Δ DIC) or change in log pseudo marginal likelihood (Δ LPML) <2 selected (Table 2).

^bCovariate estimates for each scale shown for the most parsimonious selected model via Δ DIC. Models ordered by increasing DIC. Full output in Appendix S11.

^cPositive and negative β coefficients indicate accelerated (i.e., shorter) and decelerated (i.e., longer) survival times, respectively, where $e^{-\beta}$ gives the multiplicative change in median survival time per unit covariate increase (acceleration factor). Within terms, the scale parameter, φ , describes the rate of decay in spatial correlations via $1 - e^{(-\varphi)(distance)}$, and τ^2 is the variance in the survival time in frailty (i.e., random effect).

^dCredible intervals of coefficients.

eThe test indicates one-sided hypothesis tests used to estimate evidence ratios and posterior probabilities of the statements (posterior distributions [Appendices S22-S25]).

^fThe ratio between the probability the test is true and that it is false. Evidence close to 1 indicates a low likelihood that coefficients met the test hypothesis.

gPosterior probability of the estimate.

^hReference category.

ⁱHunting period (hunting vs. nonhunting [nonhunt.]).

are already at low levels due to high anthropogenic mortality rates across the lynx's European range. We found no evidence that hunting mortality compensates for illegal killing; therefore, our results do not support the claim that hunting quotas directly help reduce illegal killing mortality (e.g., Lindsey et al., 2007).

Conservation Biology 🦄

Hunting aims to improve societal tolerance of large carnivores (Treves, 2009) rather than to change individuals' actions. Therefore, the coincidence of high illegal killing in hunted populations that we found is not necessarily unexpected. In Europe, illegal killing only appears to threaten small, reintroduced populations of lynx (Arlettaz et al., 2021; Heurich et al., 2018), whose small sizes necessitate conservation. Given the unclear relationship between illegal killing and hunting of lynx, introducing hunting management in protected populations would be highly contentious (Ghasemi, 2021). Although our results provide a first glimpse into the partially compensatory relationship between legal hunting and illegal killing, time series of tracking data in periods or areas with varying hunting quotas would be required to further disentangle them. However, these data are currently lacking. Our results imply that hunting needs to be managed carefully to ensure that the combination of natural mortality and the at least partially additive effect of hunting mortality does not exceed what the population's growth rate can tolerate. This requires effective monitoring and adaptive management feedback loops that adjust quotas (Cusack et al., 2022). Because illegal killing will likely persist, adaptive management plans, including feedback from careful monitoring, are necessary to react to dynamic situations (Andren et al., 2020). This would enable, for example, quotas to be reduced in hunted populations if illegal killing is discovered, or in hunted and protected populations, law enforcement resources could be diverted to problem areas as they emerge. Good enforcement is necessary for implementing environmental law, and in the case of large carnivores in Europe, there appears to be room for improvement (Arlettaz et al., 2021).

Roads can pose a high risk to wide-ranging species like carnivores (Bastianelli et al., 2021; Grilo et al., 2015). Vehicle collisions had a relatively low incidence compared with other sources of mortality. This is likely due to relatively low road densities where lynx often occur (e.g., Niedziałkowska et al., 2006). However, we still found that lynx that used habitats farther from roads tended to survive longer. Roads can also be dangerous for carnivores because they provide access for hunters and likewise for illegal killing (e.g., Person & Russell, 2008). Despite this, vehicle collisions must be considered an important risk that is especially concerning for small and isolated populations (Taylor et al., 2002). Vehicle collisions presented a higher risk for younger individuals, with 50% of events occurring within 3 years of age. This inevitably reduces exchange among lynx populations in Central Europe because natal dispersal is a key process for connectivity (Kramer-Schadt et al., 2004). This problem is also set to increase as road density and traffic intensity expand (Meijer et al., 2018). Improving green infrastructure and landscape permeability is a tangible intervention that may reduce lynx mortality and support population exchange. However, this depends on identifying vehicle collision hotspots for optimal placement of mitigation measures (e.g., Bil et al., 2019). Illegal killing is currently of greater magnitude and itself creates invisible barriers to lynx dispersal.

As with other large carnivores (Goodrich et al., 2008; Rauset et al., 2016), we found females had higher survival probabilities than males. This is consistent with the hypothesis that male life histories infer higher risks to secure more mates (Trivers, 1985) and is expected from sex-biased movement and dispersal (Samelius et al., 2012), with its high associated fitness costs (Lucas et al., 1994). Conversely, female lynx have higher natal philopatry (Krojerová-Prokešová et al., 2019) and appear to avoid risky behavior compared to males (Bunnefeld et al., 2006). In some hunted populations, males are at higher risk of being hunted than females because of female subquotas (Nilsen et al., 2012). In our results, male lynx in hunted populations had a higher risk of being hunted than females; however, their illegal killing rate was similar. If illegal kills occur by chance, through opportunistic encounters, poisoning, or snaring, one might expect a low sex bias, as observed. However, sexually dimorphic prey selection (e.g., Sunde & Kvam, 1997) could modulate lynx's spatial behavior and therefore risk for humancaused mortality. Sex-specific survival seems to be, therefore, the aggregation of sex-specific factors and behavior with local prey communities and human factors, such as hunting quotas. Our data set was not sufficient to consider sex-specific additive and compensatory mortality hypotheses, but these relationships could have important implications for management (e.g., Toïgo et al., 2008).

Solitary felids exhibit sex-specific seasonal activities that could affect their survival (Sandell, 2019). As expected, lynx hunting season was associated with the highest mortality risk (Andrén et al., 2022). After the hunting periods, which occur during winter, we also found winter without hunting pressure to pose a high risk. Weather conditions are unlikely to affect survival because lynx are well adapted to hunting in snow (Nilsen et al., 2009) when prey is less mobile (Mech et al., 1987). Winter could be linked to higher intraspecific competition for mates (Mattisson et al., 2013). More importantly, during winter, wildlife also leave tracks in the snow that may increase their likelihood of opportunistic illegal killing (e.g., Santiago-Ávila & Treves, 2022), especially in areas close to human activities, where detection is most likely and where prey may be aggregated (e.g., feeding sites or agricultural areas; Bunnefeld et al., 2006). Despite seasonality in lynx life history, there was little statistical evidence to differentiate between autumn (females with kittens become more mobile), spring (juveniles reach independence), and summer (kitten rearing).

Habitat selection takes place at various spatial and temporal scales to maximize fitness (Johnson, 1980). Typically, the most limiting factors for individual fitness are believed to be addressed at coarser scales (Rettie & Messier, 2000). Considering the high anthropogenic mortality rates of lynx, the hierarchical habitat selection of lynx seems consistent with the limiting factor hypothesis, wherein lynx avoid human disturbances more at the landscape scale than within the home range (Ripari et al., 2022). Habitats with little human disturbance are therefore considered an advantage for lynx occurrence (Oeser, Heurich, Kramer-Schadt, Andrén et al., 2023) and might be expected to improve a lynx's chance of survival. Following this, we hypothesized that survival depends on the use of different factors at different spatial scales. We found lynx survived longer when their home range composition and within-homerange habitat use were associated with areas with little human modification and areas farther from human infrastructure, such as roads and settlements. The evidence of this was strongest regarding habitat use at the home range scale. These results were consistent with our hypothesis based on avoidance of human disturbances at the landscape scale (H8) but did not strongly reflect the hierarchical process expected (H7) in which lynx survival was correlated more with use of habitat associated with

prey resources in the home range (H9) than with avoidance of the main mortality risk at the coarser scale. Further, we found weak to moderate evidence that use of more suitable habitat at the fine scale increased survival rates (H11).

Local studies, partly based on the same data as this study, showed disparate results. Lynx survival in Norway was shown to decrease with increasing accessibility of forests (Basille et al., 2013), whereas there was little effect of landscape on survival in neighboring Sweden (Andrén et al., 2022). In this analysis, across more study areas, lynx embedded in more humanmodified landscapes seem to be at higher risk at both fine and coarse scales. This suggests that not all lynx can minimize the fitness cost at the coarse scale and must also use habitats cautiously at the home range scale. In addition to landscape drivers, we found that some of the spatial variation and heterogeneity of risk was due to unobserved local factors. For instance, areas farther from human infrastructure may observably represent refuges; however, illegal activities can go undetected in more remote areas (Rauset et al., 2016), which would be an unobservable local risk factor. Survival is dependent on diverse local factors, which are partly difficult to describe with landscape proxies. For example, attitudes or actions opposing lynx can arise over competitive interest for game species (Treves, 2009) or livestock depredation (Abade et al., 2014). Furthermore, many processes that affect survival depend on con- and heterospecific population densities, including prey availability, competitive killing, or disease transmission (Brøseth et al., 2010; Murray et al., 1999; Palomares & Caro, 1999). Although some factors are ultimately conveyed by habitat use (e.g., individuals forced into less suitable areas due to high conspecific density), not all factors can be quantified, such as prey vulnerability, human activities, or cultural aspects of tolerance, and especially the behaviors of small groups of humans who engage in illegal killing. By statistically accounting for the variability of these confounding factors, we could understand the effect of habitat use alone. Our results thereby corroborate the prevailing wisdom that conserving habitat and preventing further fragmentation are fundamental ingredients for lynx conservation.

We showed that lynx survival was partly driven by landscape factors, particularly correlates of human influences. Habitats with little human disturbance and human infrastructure offer lynx the best survival chances, provided unobserved factors are not detrimental. The at best partially compensatory relationship between anthropogenic and other mortality causes highlights that Europe's lynx occurrences are at high risk of mortality. In fact, with natural mortality already at low levels in hunted and protected populations, mortality causes are likely close to additive already. For this reason, the persistence of small lynx populations should not be taken for granted. Whether lynx can thrive in novel patches will depend on dynamic local conditions, including societal perceptions, management actions (e.g., assisted dispersal), and ecological connectivity among patches (Jaroszewicz et al., 2021; Linnell et al., 2015). These factors should be considered in future research to better understand spatial variation in population dynamics of lynx.

AFFILIATIONS

¹Chair of Wildlife Ecology and Management, Albert Ludwigs University of Freiburg, Freiburg, Germany

Conservation Biology

²Department of National Park Monitoring and Animal Management, Bavarian Forest National Park, Grafenau, Germany

³Leibniz Institute for Zoo and Wildlife Research (IZW), Berlin, Germany

⁴Geography Department, Humboldt-Universität zu Berlin, Berlin, Germany

⁵Luchsprojekt Harz, Nationalparkverwaltung Harz, Außenstelle Oderhaus, Sankt Andreasberg, Germany

⁶Grimsö Wildlife Research Station, Department of Ecology, Swedish University of Agricultural Sciences, Riddarhyttan, Sweden

⁷Latvian State Forest Research Institute Silava, Salaspils, Latvia

⁸Faculty of Forestry and Wood Sciences, Czech University of Life Sciences, Prague, Czech Republic

⁹Department of Research and Nature Protection, Šumava National Park Administration, Kašperské Hory, Czech Republic

¹⁰Stiftung KORA, Muri bei Bern, Switzerland

¹¹Slovenia Forest Service, Ljubljana, Slovenia

¹²Direction de la Recherche et de l'Appui Scientifique, Office Français de la Biodiversité, Gières, France

¹³Department of Forest Ecology, Faculty of Forestry and Wood Technology, Mendel University in Brno, Brno, Czech Republic

¹⁴Luchsbeauftragter, Nationalpark Kalkalpen, Nationalpark Zentrum Molln, Molln, Austria

¹⁵Forstliche Versuchs- und Forschungsanstalt, Baden-Württemberg (FVA), Freiburg, Germany

¹⁶Biotechnical Faculty, University of Ljubljana, Ljubljana, Slovenia

¹⁷Centro de Ecología, Instituto Venezolano de Investigaciones Científicas (IVIC), Caracas, Venezuela

¹⁸Mammal Research Institute, Polish Academy of Sciences, Białowieża, Poland

¹⁹Department of Zoology, Institute of Ecology and Earth Sciences, University of Tartu, Tartu, Estonia

²⁰Institute of Vertebrate Biology, Czech Academy of Sciences, Brno, Czech Republic

²¹Department of Zoology, Fisheries, Hydrobiology and Apiculture, Faculty of Agri-Sciences, Mendel University in Brno, Brno, Czech Republic

²²Department of Applied Zoology and Wildlife Management, Faculty of Forestry, Technical University in Zvolen, Zvolen, Slovakia

²³DIANA – Carpathian Wildlife Research, Banská Bystrica, Slovakia

²⁴Department of Biology, Faculty of Veterinary Medicine, University of Zagreb, Zagreb, Croatia

²⁵Carnivore Conservation Programme, Friends of the Earth Czech Republic, Olomouc, Czech Republic

²⁶Norwegian Institute for Nature Research, Lillehammer, Norway

²⁷Department of Forestry and Wildlife Management, Campus Evenstad, Inland Norway University for Applied Science, Koppang, Norway

²⁸Norwegian Institute for Nature Research, Trondheim, Norway

²⁹Macedonian Ecological Society, Skopje, North Macedonia

³⁰Progetto Lince Italia, Tarvisio, Italy

³¹Norwegian Institute for Nature Research, Oslo, Norway

³²Institute of Nature Conservation, Polish Academy of Sciences, Krakow, Poland

³³Department of Wildlife Management and Nature Protection, Karlovac University of Applied Sciences, Karlovac, Croatia

³⁴National Zoological Garden Bojnice, Bojnice, Slovakia

³⁵Department of Ecology and Evolution, University of Lausanne, Lausanne, Switzerland

³⁶Chair of Applied Animal Ecology, Institute of Ecology, Technische Universität Berlin, Berlin, Germany

ACKNOWLEDGMENTS

We especially thank the game wardens, wildlife veterinarians, field personnel, interns, civil servants, students, and hunters in the research sites for their essential help with capturing and monitoring of lynx. This paper was conceived and written within the EUROLYNX collaborative project (paper 06 of the EUROLYNX series; www.eurolynx.org). Open access funding was enabled and organized by Projekt DEAL.

ORCID

J. Premier https://orcid.org/0000-0003-3225-6038 M. L. Bastianelli https://orcid.org/0000-0003-2309-4903 J. Oeser https://orcid.org/0000-0003-3216-6817 O. Anders https://orcid.org/0000-0002-0505-5743 H. Andren https://orcid.org/0000-0002-5616-2426 M. Aronsson https://orcid.org/0000-0002-9026-3765 G. Bagrade https://orcid.org/0000-0002-1031-0665 E. Belotti https://orcid.org/0000-0002-6072-003X C. Breitenmoser-Würsten https://orcid.org/0000-0002-5228-2957

L. Bufka https://orcid.org/0000-0001-5550-7712 R. Černe https://orcid.org/0000-0003-4111-4066 J. Červený https://orcid.org/0000-0002-2723-0936 N. Drouet-Hoguet https://orcid.org/0000-0002-2610-6917 M. Ďula https://orcid.org/0000-0003-2675-4575 L. Hočevar https://orcid.org/0000-0002-2346-1390 W. Jędrzejewski https://orcid.org/0000-0002-9934-0683 R. Kowalczyk https://orcid.org/0000-0001-7264-2158 M. Krofel https://orcid.org/0000-0002-2010-5219 J. Krojerová-Prokešová https://orcid.org/0000-0002-0067-6196

J. Kubala https://orcid.org/0000-0003-2051-508X *J. Kusak* b https://orcid.org/0000-0002-5544-6486 *M. Kutal* https://orcid.org/0000-0003-3857-5419 I.D. C. Linnell https://orcid.org/0000-0002-8370-5633 J. Mattisson D https://orcid.org/0000-0001-6032-5307 D. Melovski b https://orcid.org/0000-0002-1072-6978 A. Molinari-Jobin b https://orcid.org/0000-0002-3729-2532 J. Odden b https://orcid.org/0000-0002-6275-8648 H. Okarma b https://orcid.org/0000-0002-8912-3546 A. Ornicāns b https://orcid.org/0000-0002-1478-3080 N. Pagon b https://orcid.org/0009-0001-2462-5203 *J. Persson* b https://orcid.org/0000-0003-1405-7561 K. Schmidt b https://orcid.org/0000-0002-9043-291X M. Sindičić https://orcid.org/0000-0002-1391-3648 F. Zimmermann b https://orcid.org/0000-0003-3861-246X S. Kramer-Schadt b https://orcid.org/0000-0002-9269-4446 *M. Heurich* https://orcid.org/0000-0003-0051-2930

REFERENCES

- Abade, L., Macdonald, D. W., & Dickman, A. J. (2014). Assessing the relative importance of landscape and husbandry factors in determining large carnivore depredation risk in Tanzania's Ruaha landscape. *Biological Conservation*, 180, 241–248.
- Andrén, H., Aronsson, M., López-Bao, J. V., Samelius, G., Chapron, G., Rauset, G. R., Hemmingmoore, H., & Persson, J. (2022). Season rather than habitat affects lynx survival and risk of mortality in the human-dominated landscape of southern Sweden. *Wildlife Biology*, 2022(1), Article e01008.

- Andrén, H., Hobbs, N. T., Aronsson, M., Brøseth, H., Chapron, G., Linnell, J. D., Odden, J., Persson, J., & Nilsen, E. B. (2020). Harvest models of small populations of a large carnivore using Bayesian forecasting. *Ecological Applications*, 30(3), Article e02063.
- Andrén, H., Linnell, J. D., Liberg, O., Andersen, R., Danell, A., Karlsson, J., Odden, J., Moa, P. F., Ahlqvist, P., Kvam, T., Franzén, R., & Segerström, P. (2006). Survival rates and causes of mortality in Eurasian lynx (*Lynx lynx*) in multi-use landscapes. *Biological Conservation*, 131(1), 23–32.
- Arlettaz, R., Chapron, G., Kéry, M., Klaus, E., Mettaz, S., Roder, S., Vignali, S., Zimmermann, F., & Braunisch, V. (2021). Poaching threatens the establishment of a lynx population, highlighting the need for a centralized judiciary approach. *Frontiers in Conservation Science*, 2, Article 665000.
- Basille, M., Van Moorter, B., Herfindal, I., Martin, J., Linnell, J. D., Odden, J., Andersen, R., & Gaillard, J. M. (2013). Selecting habitat to survive: The impact of road density on survival in a large carnivore. *PLoS ONE*, 8(7), Article e65493.
- Bastianelli, M. L., Premier, J., Herrmann, M., Anile, S., Monterroso, P., Kuemmerle, T., Dormann, C. F., Streif, S., Jerosch, S., Götz, M., Simon, O., Moleón, M., Gil-Sánchez, J. M., Biró, Z., Dekker, J., Severon, A., Krannich, A., Hupe, K., Germain, E., ... Heurich, M. (2021). Survival and causespecific mortality of European wildcat (*Felis silvestris*) across Europe. *Biological Conservation*, 261, Article 109239.
- Benson, J. F., Dougherty, K. D., Beier, P., Boyce, W. M., Cristescu, B., Gammons, D. J., Garcelon, D. K., Higley, J. M., Martins, Q. E., Nisi, A. C., Riley, S. P. D., Sikich, J. A., Stephenson, T. R., Vickers, T. W., Wengert, G. M., Wilmers, C. C., Wittmer, H. U., & Dellinger, J. A. (2023). The ecology of human-caused mortality for a protected large carnivore. *Proceedings of the National Academy of Sciences of the United States of America*, 120(13), Article e2220030120.
- Bíl, M., Andrášik, R., Duľa, M., & Sedoník, J. (2019). On reliable identification of factors influencing wildlife-vehicle collisions along roads. *Journal of Environmental Management*, 237, 297–304.
- Boyce, M. S., Sinclair, A. R. E., & White, G. C. (1999). Seasonal compensation of predation and harvesting. *Oikas*, 87, 419–426.
- Bradburn, M. J., Clark, T. G., Love, S. B., & Altman, D. G. (2003). Survival analysis part II: Multivariate data analysis—An introduction to concepts and methods. *British Journal of Cancer*, 89(3), 431–436.
- Breitenmoser-Würsten, C., Vandel, J. M., Zimmermann, F., & Breitenmoser, U. (2007). Demography of lynx *Lynx lynx* in the Jura Mountains. *Wildlife Biology*, 13(4), 381–392.
- Brodie, J., Johnson, H., Mitchell, M., Zager, P., Proffitt, K., Hebblewhite, M., Kauffman, M., Johnson, B., Bissonette, J., Bishop, C., Gude, J., Herbert, J., Hersey, K., Hurley, M., Lukacs, P. M., McCorquodale, S., McIntire, E., Nowak, J., Sawyer, H., ... White, P. J. (2013). Relative influence of human harvest, carnivores, and weather on adult female elk survival across western North America. *Journal of Applied Ecology*, 50(2), 295–305.
- Brøseth, H., Flagstad, Ø., Wärdig, C., Johansson, M., & Ellegren, H. (2010). Large-scale noninvasive genetic monitoring of wolverines using scats reveals density dependent adult survival. *Biological Conservation*, 143(1), 113–120.
- Bunnefeld, N., Linnell, J. D., Odden, J., Van Duijn, M. A. J., & Andersen, R. (2006). Risk taking by Eurasian lynx (*Lynx lynx*) in a human-dominated landscape: Effects of sex and reproductive status. *Journal of Zoology*, 270(1), 31–39.
- Burham, K. P., & Anderson, D. R. (1984). Tests of compensatory vs. additive hypotheses of mortality in mallards. *Ecology*, 65(1), 105–112.
- Bürkner, P. C. (2017). brms: An R package for Bayesian multilevel models using Stan. *Journal of Statistical Software*, 80, 1–28.
- Carbone, C., Cowlishaw, G., Isaac, N. J., & Rowcliffe, J. M. (2005). How far do animals go? Determinants of day range in mammals. *The American Naturalist*, 165(2), 290–297.
- Carter, N. H., López-Bao, J. V., Bruskotter, J. T., Gore, M., Chapron, G., Johnson, A., Epstein, Y., Shrestha, M., Frank, J., Ohrens, O., & Treves, A. (2017). A conceptual framework for understanding illegal killing of large carnivores. *Ambio*, 46(3), 251–264.
- Cribari-Neto, F., & Zeileis, A. (2010). Beta regression in R. Journal of Statistical Software, 34, 1–24.
- Cusack, J. J., Nilsen, E. B., Israelsen, M. F., Andrén, H., Grainger, M., Linnell, J. D., Odden, J., & Bunnefeld, N. (2022). Quantifying the checks and balances

of collaborative governance systems for adaptive carnivore management. *Journal of Applied Ecology*, 59(4), 1038–1049.

Conservation Biology 🚿

- DeCesare, N. J., Hebblewhite, M., Bradley, M., Hervieux, D., Neufeld, L., & Musiani, M. (2014). Linking habitat selection and predation risk to spatial variation in survival. *Journal of Animal Ecology*, 83(2), 343– 352.
- de Wreede, L. C., Fiocco, M., & Putter, H. (2011). mstate: An R package for the analysis of competing risks and multi-state models. *Journal of Statistical Software*, 38(7), 1–30.
- Errington, P. L. (1956). Factors limiting higher vertebrate populations. Science, 124(3216), 304–307.
- Fahrig, L. (2007). Non-optimal animal movement in human-altered landscapes. *Functional Ecology*, 21(6), 1003–1015.
- Fieberg, J., & DelGiudice, G. D. (2009). What time is it? Choice of time origin and scale in extended proportional hazards models. *Ecology*, 90(6), 1687–1697.
- Filla, M., Premier, J., Magg, N., Dupke, C., Khorozyan, I., Waltert, M., Bufka, L., & Heurich, M. (2017). Habitat selection by Eurasian lynx (*Lynx lynx*) is primarily driven by avoidance of human activity during day and prey availability during night. *Ecology and Evolution*, 7(16), 6367–6381.
- Gaillard, J. M., Hebblewhite, M., Loison, A., Fuller, M., Powell, R., Basille, M., & Van Moorter, B. (2010). Habitat–performance relationships: Finding the right metric at a given spatial scale. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 365(1550), 2255–2265.
- Geskus, R. B. (2011). Cause-specific cumulative incidence estimation and the fine and gray model under both left truncation and right censoring. *Biometrics*, 67(1), 39–49.
- Ghasemi, B. (2021). Trophy hunting and conservation: Do the major ethical theories converge in opposition to trophy hunting? *People and Nature*, 3(1), 77–87.
- Gilroy, J. J., Ordiz, A., & Bischof, R. (2015). Carnivore coexistence: Value the wilderness. Science, 347(6220), 382.
- Goodrich, J. M., Kerley, L. L., Smirnov, E. N., Miquelle, D. G., McDonald, L., Quigley, H. B., Hornocker, M. G., & McDonald, T. (2008). Survival rates and causes of mortality of Amur tigers on and near the Sikhote-Alin Biosphere Zapovednik. *Journal of Zoology*, 276(4), 323–329.
- Grilo, C., Smith, D. J., & Klar, N. (2015). Carnivores: Struggling for survival in roaded landscapes. In R. van der Ree, D. J. Smith, & C. Grilo (Eds.), *Handbook* of road ecology (pp. 300–312). John Wiley & Sons.
- Halstead, B. J., Wylie, G. D., Coates, P. S., Valcarcel, P., & Casazza, M. L. (2012). Bayesian shared frailty models for regional inference about wildlife survival. *Animal Conservation*, 15(2), 117–124.
- Heisey, D. M., & Fuller, T. K. (1985). Evaluation of survival and cause-specific mortality rates using telemetry data. *The Journal of Wildlife Management*, 49(3), 668–674.
- Heurich, M., Schultze-Naumburg, J., Piacenza, N., Magg, N., Červený, J., Engleder, T., Herdtfelder, M., Sladova, M., & Kramer-Schadt, S. (2018). Illegal hunting as a major driver of the source-sink dynamics of a reintroduced lynx population in Central Europe. *Biological Conservation*, 224, 355–365.
- Jaroszewicz, B., Nowak, K., & Zmihorski, M. (2021). Poland's border wall threatens ancient forest. *Science*, 374(6571), 1063–1063.
- Johnson, D. H. (1980). The comparison of usage and availability measurements for evaluating resource preference. *Ecology*, 61(1), 65–71.
- Kaplan, E. L., & Meier, P. (1958). Nonparametric estimation from incomplete observations. *Journal of the American Statistical Association*, 53(282), 457–481.
- Kassambara, A., Kosinski, M., Biecek, P., & Fabian, S. (2017). Package 'survminer': Drawing survival curves using 'ggplot2'. R package version 0.31.
- Koons, D. N., Gamelon, M., Gaillard, J. M., Aubry, L. M., Rockwell, R. F., Klein, F., Choquet, R., & Gimenez, O. (2014). Methods for studying cause-specific senescence in the wild. *Methods in Ecology and Evolution*, 5(9), 924–933.
- Kramer-Schadt, S., Revilla, E., Wiegand, T., & Breitenmoser, U. (2004). Fragmented landscapes, road mortality and patch connectivity: Modelling influences on the dispersal of Eurasian lynx. *Journal of Applied Ecology*, 41(4), 711–723.
- Krofel, M., Treves, A., Ripple, W. J., Chapron, G., & López-Bao, J. V. (2015). Hunted carnivores at outsized risk. *Science*, 350(6260), 518–519.
- Krojerová-Prokešová, J., Turbaková, B., Jelenčič, M., Bojda, M., Kutal, M., Skrbinšek, T., Koubek, P., & Bryja, J. (2019). Genetic constraints of pop-

and Conditions (https://onlinelibrary.wiley.com

onditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons

16 of 17

Conservation Biology 🗞

ulation expansion of the Carpathian lynx at the western edge of its native distribution range in Central Europe. *Heredity*, 122(6), 785–799.

- Lamarca, R., Alonso, J., Gomez, G., & Muñoz, Á. (1998). Left-truncated data with age as time scale: An alternative for survival analysis in the elderly population. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, 53(5), M337–M343.
- Lamb, C. T., Ford, A. T., McLellan, B. N., Proctor, M. F., Mowat, G., Ciarniello, L., Nielsen, S. E., & Boutin, S. (2020). The ecology of human–carnivore coexistence. *Proceedings of the National Academy of Sciences of the United States of America*, 117(30), 17876–17883.
- Lindsey, P. A., Roulet, P. A., & Romañach, S. S. (2007). Economic and conservation significance of the trophy hunting industry in sub-Saharan Africa. *Biological Conservation*, 134(4), 455–469.
- Linnell, J. D., Brøseth, H., Odden, J., & Nilsen, E. B. (2010). Sustainably harvesting a large carnivore? Development of Eurasian lynx populations in Norway during 160 years of shifting policy. *Environmental Management*, 45(5), 1142–1154.
- Linnell, J. D., Kaczensky, P., Wotschikowsky, U., Lescureux, N., & Boitani, L. (2015). Framing the relationship between people and nature in the context of European conservation. *Conservation Biology*, 29(4), 978–985.
- Lucas, J. R., Waser, P. M., & Creel, S. R. (1994). Death and disappearance: Estimating mortality risks associated with philopatry and dispersal. *Behavioral Ecology*, 5(2), 135–141.
- Marboutin, E., Bray, Y., Péroux, R., Mauvy, B., & Lartiges, A. (2003). Population dynamics in European hare: Breeding parameters and sustainable harvest rates. *Journal of Applied Ecology*, 40(3), 580–591.
- Marti, I., & Ryser-Degiorgis, M. P. (2018). A tooth wear scoring scheme for age estimation of the Eurasian lynx (*Lynx lynx*) under field conditions. *European Journal of Wildlife Research*, 64(4), Article 37.
- Mattisson, J., Linnell, J. D., Anders, O., Belotti, E., Breitenmoser-Würsten, C., Bufka, L., Fuxjäger, C., Heurich, M., Ivanov, G., Jędrzejewski, W., Kont, R., Kowalczyk, R., Krofel, M., Melovski, D., Mengüllüoğlu, D., Middelhoff, T. L., Molinari-Jobin, A., Odden, J., Ozoliņš, J., ... Andrén, H. (2022). Timing and synchrony of birth in Eurasian lynx across Europe. *Ecology and Evolution*, 12(8), Article e9147.
- Mattisson, J., Odden, J., Linnell, J. D., Painer, J., Persson, J., & Andrén, H. (2020). Parturition dates in wild Eurasian lynx: Evidence of a second oestrus? *Mammalian Biology*, 100, 549–552.
- Mattisson, J., Segerström, P., Persson, J., Aronsson, M., Rauset, G. R., Samelius, G., & Andrén, H. (2013). Lethal male-male interactions in Eurasian lynx. *Mammalian Biology*, 78, 304–308.
- Mech, L. D., McRoberts, R. E., Peterson, R. O., & Page, R. E. (1987). Relationship of deer and moose populations to previous winters' snow. *The Journal of Animal Ecology*, 56(2), 615–627.
- Medellín, R. A. (1999). Sustainable harvest for conservation. *Conservation Biology*, 13(2), 225–227.
- Meijer, J. R., Huijbregts, M. A., Schotten, K. C., & Schipper, A. M. (2018). Global patterns of current and future road infrastructure. *Environmental Research Letters*, 13(6), Article 064006.
- Moss, W. E., Alldredge, M. W., & Pauli, J. N. (2016). Quantifying risk and resource use for a large carnivore in an expanding urban-wildland interface. *Journal of Applied Ecology*, 53(2), 371–378.
- Murray, D. L., Kapke, C. A., Evermann, J. F., & Fuller, T. K. (1999). Infectious disease and the conservation of free-ranging large carnivores. *Animal Conservation*, 2(4), 241–254.
- Murray, D. L., & Patterson, B. R. (2006). Wildlife survival estimation: Recent advances and future directions. *Journal of Wildlife Management*, 70(6), 1499– 1503.
- Murray, D. L., Smith, D. W., Bangs, E. E., Mack, C., Oakleaf, J. K., Fontaine, J., Boyd, D., Jiminez, M., Niemeyer, C., Meier, T. J., Stahler, D., Holyan, J., & Asher, V. J. (2010). Death from anthropogenic causes is partially compensatory in recovering wolf populations. *Biological Conservation*, 143(11), 2514–2524.
- Naef-Daenzer, B., Korner-Nievergelt, F., Fiedler, W., & Grüebler, M. U. (2017). Bias in ring-recovery studies: Causes of mortality of little owls *Athene noctua* and implications for population assessment. *Journal Avian Biology*, 48(2), 266– 274.

- Niedziałkowska, M., Jędrzejewski, W., Mysłajek, R. W., Nowak, S., Jędrzejewska, B., & Schmidt, K. (2006). Environmental correlates of Eurasian lynx occurrence in Poland—Large scale census and GIS mapping. *Biological Conservation*, 133(1), 63–69.
- Nilsen, E. B., Brøseth, H., Odden, J., & Linnell, J. D. (2012). Quota hunting of Eurasian lynx in Norway: Patterns of hunter selection, hunter efficiency and monitoring accuracy. *European Journal of Wildlife Research*, 58(1), 325– 333.
- Nilsen, E. B., Linnell, J. D., Odden, J., & Andersen, R. (2009). Climate, season, and social status modulate the functional response of an efficient stalking predator: The Eurasian lynx. *Journal of Animal Ecology*, 78(4), 741–751.
- Nisi, A. C., Benson, J. F., King, R., & Wilmers, C. C. (2023). Habitat fragmentation reduces survival and drives source–sink dynamics for a large carnivore. *Ecological Applications*, 33(4), Article e2822.
- Oeser, J., Heurich, M., Kramer-Schadt, S., Andrén, H., Bagrade, G., Belotti, E., Bufka, L., Breitenmoser-Würsten, C., Černe, R., Dula, M., Fuxjäger, C., Gomerčić, T., Jędrzejewski, W., Kont, R., Koubek, P., Kowalczyk, R., Krofel, M., Krojerová-Prokešová, J., Kubala, J., ... Kuemmerle, T. (2023). Prerequisites for coexistence: Human pressure and refuge habitat availability shape continental-scale habitat use patterns of a large carnivore. *Landscape Ecology*, 38(7), 1713–1728.
- Oeser, J., Heurich, M., Kramer-Schadt, S., Mattisson, J., Krofel, M., Krojerová-Prokešová, J., Zimmermann, F., Anders, O., Andrén, H., Bagrade, G., Belotti, E., Breitenmoser-Würsten, C., Bufka, L., Černe, R., Drouet-Hoguet, N., Duľa, M., Fuxjäger, C., Gomerčić, T., Jędrzejewski, W., ... Kuemmerle, T. (2023). Integrating animal tracking datasets at a continental scale for mapping Eurasian lynx habitat. *Diversity and Distributions*, 29(12), 1546–1560.
- Oriol-Cotterill, A., Valeix, M., Frank, L. G., Riginos, C., & Macdonald, D. W. (2015). Landscapes of coexistence for terrestrial carnivores: The ecological consequences of being downgraded from ultimate to penultimate predator by humans. *Oikos*, 124(10), 1263–1273.
- Palomares, F., & Caro, T. M. (1999). Interspecific killing among mammalian carnivores. *The American Naturalist*, 153(5), 492–508.
- Person, D. K., & Russell, A. L. (2008). Correlates of mortality in an exploited wolf population. *The Journal of Wildlife Management*, 72(7), 1540–1549.
- R Core Team. (2022). R: A language and environment for statistical computing. R Foundation for Statistical Computing, https://www.R-project.org/
- Rauset, G. R., Andrén, H., Swenson, J. E., Samelius, G., Segerström, P., Zedrosser, A., & Persson, J. (2016). National parks in northern Sweden as refuges for illegal killing of large carnivores. *Conservation Letters*, 9(5), 334–341.
- Redpath, S. M., Young, J., Evely, A., Adams, W. M., Sutherland, W. J., Whitehouse, A., Amar, A., Lambert, R. A., Linnell, J. D., Watt, A., & Gutierrez, R. J. (2013). Understanding and managing conservation conflicts. *Trends in Ecology & Evolution*, 28(2), 100–109.
- Rettie, W. J., & Messier, F. (2000). Hierarchical habitat selection by woodland caribou: Its relationship to limiting factors. *Eugraphy*, 23(4), 466–478.
- Ripari, L., Premier, J., Belotti, E., Bluhm, H., Breitenmoser-Würsten, C., Bufka, L., Červený, J., Drouet-Hoguet, N., Fuxjäger, C., Jędrzejewski, W., Kont, R., Koubek, P., Kowalczyk, R., Krofel, M., Krojerová-Prokešová, J., Molinari-Jobin, A., Okarma, H., Oliveira, T., Remm, J., ... Heurich, M. (2022). Human disturbance is the most limiting factor driving habitat selection of a large carnivore throughout Continental Europe. *Biological Conservation*, 266, Article 109446.
- Samelius, G., Andrén, H., Liberg, O., Linnell, J. D. C., Odden, J., Ahlqvist, P., Segerström, P., & Sköld, K. (2012). Spatial and temporal variation in natal dispersal by Eurasian lynx in Scandinavia. *Journal of Zoology*, 286(2), 120–130.
- Sandell, M. (2019). The mating tactics and spacing patterns of solitary carnivores. In J. L. Gittleman (Ed.), *Carnivore behavior, ecology, and evolution* (pp. 164–182). Cornell University Press.
- Sandercock, B. K., Nilsen, E. B., Brøseth, H., & Pedersen, H. C. (2011). Is hunting mortality additive or compensatory to natural mortality? Effects of experimental harvest on the survival and cause-specific mortality of willow ptarmigan. *Journal of Animal Ecology*, 80(1), 244–258.
- Santiago-Ávila, F. J., & Treves, A. (2022). Poaching of protected wolves fluctuated seasonally and with non-wolf hunting. *Scientific Reports*, 12(1), Article 1738.

- Saracco, J. F., Royle, J. A., DeSante, D. F., & Gardner, B. (2010). Modeling spatial variation in avian survival and residency probabilities. *Ecology*, 91(7), 1885– 1891.
- Schmidt-Posthaus, H., Breitenmoser-Würsten, C., Posthaus, H., Bacciarini, L., & Breitenmoser, U. (2002). Causes of mortality in reintroduced Eurasian lynx in Switzerland. *Journal of Wildlife Diseases*, 38(1), 84–92.
- Signer, J., Fieberg, J., & Avgar, T. (2019). Animal movement tools (amt): R package for managing tracking data and conducting habitat selection analyses. *Ecology and Evolution*, 9(2), 880–890.
- Sih, A. (2013). Understanding variation in behavioural responses to humaninduced rapid environmental change: A conceptual overview. *Animal Behaviour*, 85(5), 1077–1088.
- Sindičić, M., Gomerčić, T., Kusak, J., Slijepčević, V., Huber, Đ., & Frković, A. (2016). Mortality in the Eurasian lynx population in Croatia over the course of 40 years. *Mammalian Biology*, 81, 290–294.
- Smith, D. W., Bangs, E. E., Oakleaf, J. K., Mack, C., Fontaine, J., Boyd, D., Jimenez, M., Pletscher, D. H., Niemeyer, C. C., Meier, T. J., Stahler, D. R., Holyan, J., Asher, V. J., & Murray, D. L. (2010). Survival of colonizing wolves in the northern Rocky Mountains of the United States, 1982–2004. *The Journal of Wildlife Management*, 74(4), 620–634.
- Sunde, P., & Kvam, T. (1997). Diet patterns of Eurasian lynx Lynx lynx: what causes sexually determined prey size segregation? *Acta Theriologica*, 42, 189– 201.
- Suraci, J. P., Frank, L. G., Oriol-Cotterill, A., Ekwanga, S., Williams, T. M., & Wilmers, C. C. (2019). Behavior-specific habitat selection by African lions may promote their persistence in a human-dominated landscape. *Ecology*, 100(4), Article e02644.
- Suutarinen, J., & Kojola, I. (2017). Poaching regulates the legally hunted wolf population in Finland. *Biological Conservation*, 215, 11–18.
- Taylor, S. K., Buergelt, C. D., Roelke-Parker, M. E., Homer, B. L., & Rotstein, D. S. (2002). Causes of mortality of free-ranging Florida panthers. *Journal of Wildlife Diseases*, 38(1), 107–114.
- Therneau, T. (2021). A package for survival analysis in R. R package version 3.2-13. https://CRAN.R-project.org/package=survival
- Toigo, C., Servanty, S., Gaillard, J. M., Brandt, S., & Baubet, E. (2008). Disentangling natural from hunting mortality in an intensively hunted wild boar population. *The Journal of Wildlife Management*, 72(7), 1532–1539.
- Treves, A. (2009). Hunting for large carnivore conservation. Journal of Applied Ecology, 46(6), 1350–1356.
- Trivers, R. (1985). Social evolution. Benjamin/Cumming.
- Weingarth, K., Heibl, C., Knauer, F., Zimmermann, F., Bufka, L., & Heurich, M. (2012). First estimation of Eurasian lynx (Lynx lynx) abundance and density using digital cameras and capture–recapture techniques in a German national park. *Animal Biodiversity and Conservation*, 35(2), 197–207.

Conservation Biology 🔌 📋 17 of 17

- Weingarth, K., Zeppenfeld, T., Heibl, C., Heurich, M., Bufka, L., Daniszová, K., & Müller, J. (2015). Hide and seek: Extended camera-trap session lengths and autumn provide best parameters for estimating lynx densities in mountainous areas. *Biodiversity and Conservation*, 24, 2935–2952.
- Wickham, H. (2016). ggplot2: Elegant graphics for data analysis. Springer-Verlag.
- Wolfe, M. L., Koons, D. N., Stoner, D. C., Terletzky, P., Gese, E. M., Choate, D. M., & Aubry, L. M. (2015). Is anthropogenic cougar mortality compensated by changes in natural mortality in Utah? Insight from long-term studies. *Biological Conservation*, 182, 187–196.
- Zabor, E. C., & Gonen, M. (2019). condsurv: Conditional survival estimates and plots. R package version 1.0.0. http://www.emilyzabor.com/condsurv/
- Zhou, H., & Hanson, T. (2015). Bayesian spatial survival models. In R. Mitra & P. Müller (Eds.), *Nonparametric Bayesian inference in biostatistics* (pp. 215–246). Springer.
- Zhou, H., & Hanson, T. (2018). A unified framework for fitting Bayesian semiparametric models to arbitrarily censored survival data, including spatially referenced data. *Journal of the American Statistical Association*, 113(522), 571–581.
- Zhou, H., Hanson, T., & Zhang, J. (2020). spBayesSurv: Fitting Bayesian spatial survival models using R. *Journal of Statistical Software*, 92, 1–33.
- Zimmermann, F., Breitenmoser-Würsten, C., & Breitenmoser, U. (2005). Natal dispersal of Eurasian lynx (*Lynx lynx*) in Switzerland. *Journal of Zoology*, 267(4), 381–395.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Premier, J., Bastianelli, M. L., Oeser, J., Anders, O., Andren, H., Aronsson, M., Bagrade, G., Belotti, E., Breitenmoser-Würsten, C., Bufka, L., Černe, R., Červený, J., Drouet-Hoguet, N., Ďuľa, M., Fuxjäger, C., Herdtfelder, M., Hočevar, L., Jędrzejewski, W., Kont, R., ... Heurich, M. (2025). Survival of Eurasian lynx in the human-dominated landscape of Europe. *Conservation Biology*, e14439. https://doi.org/10.1111/cobi.14439